

More records of North American vascular plants from the Sperry Herbarium

NEIL SNOW¹, RACHEL STYERS-OSBORN², RYLAN MASON², ALAINA FULKERSON², GRACE GIBBS², JORDAN HAWORTH², CHRISTINE HELMAN², SYDNEY HULVEY², GABE MCCLAIN², ISABELLE MCMAHON², ALAENAH MICHAUD², GAGE SINGER², AND KHLOEY STRINGER²

ABSTRACT. — *Heliotropium lasiocarpum* is reported new for Oklahoma; this is the second report from North America. Sixty-five additional new county records of vascular plants are reported from 26 counties in Kansas (7 taxa), Missouri (41 taxa) and Oklahoma (17 taxa). Some taxa are reported as new from multiple counties. Of the 65 total records, 40 are native and 25 are nonnative to North America. These records derive primarily from recent graduate projects, floristic studies, and class activities at Pittsburg State University from 2022–2024. We discuss the unexpected proliferation of county records regionally and the implications of this level of documentation.

INTRODUCTION

Important tasks in plant taxonomy include formal descriptions of new plant taxa, updating taxonomic knowledge with revisions and monographs, producing checklists and Floras, and inferring evolutionary relationships among taxa. An equally important but sometimes less acknowledged goal is documenting the geographical distributions of taxa with vouchered herbarium specimens.

Increasing evidence indicates that our knowledge of the distributions of vascular plants in North America remains far from complete at the county level. Over the past eleven field seasons (2014–2024), work at the Sperry Herbarium (KSP) at Pittsburg State University (PSU) has documented with voucher specimens over 700 new records of new vascular plants at the state or county level (Snow, in prep.). Although many records from KSP have been published (e.g., Snow 2017, Snow et al. 2017, Pryer et al. 2019, Hammesfahr et al. 2020, Daines et al. 2022), some were reported directly to the Biota of North America Program (Kartesz 2024) without formal publication.

These new records derive from several sources, including required plant collections from courses at PSU such as Taxonomy of Vascular Plants, Grass Taxonomy, and Wetland Plants; these specimens are typically within a 300 km radius of Pittsburg, Kansas. Others came from a master's thesis (Styers 2022) and an ongoing floristic survey of Bates County, Missouri. The remaining records are from opportunistic collections of the first author or from existing herbarium specimens at KSP that, upon digitization, appeared to be new records. Most of the records reported below were collected between 2022 and 2024.

¹ NEIL SNOW (corresponding author) — T.M. Sperry Herbarium, Department of Biology, Pittsburg State University, Pittsburg, KS 66762. email: nsnow@pittstate.edu.

² Department of Biology, Pittsburg State University, Pittsburg, KS 66762.

METHODS

Plants were collected following standard fieldwork practices for temperate climates, wherein specimens are pressed immediately in the field in presses with cardboard ventilators and later dried over artificial heat, typically for three to five days.

To the best of our knowledge, each record represents the first vouchered occurrence for the indicated county or state based on SEINet (2025) and BONAP (Kartesz 2024) through 1 April 2025. *Heliotropium lasiocarpum* is new to Oklahoma; all other reports represent county records.

Sources used to identify the taxa reported relied primarily on Yatskievych (1999, 2006, 2013) but also included Haddock et al. (2015), Stubbendieck et al. (2017), Brummitt (1972) and the treatment of Cyperaceae (by various authors) in *Flora of North America* (Volume 26). Common names mostly reflect those found in the volumes of Yatskievych for Missouri, Stubbendieck et al. (2017) for Poaceae, and BONAP (Kartesz 2024). Authors were responsible for identifying specimens collected under their names and confirming data in DarwinCore spreadsheets (Wieczorek et al. 2012); the lead author later confirmed each determination. All digital data associated with the reported specimens are available from the Consortium of Northern Great Plains Herbaria via SEINet (2025). Given the pending release of an updated online version of BONAP (J. Kartesz, pers. comm., Dec. 2024), these records (taxa + collector names + numbers + KSP barcodes) also have been submitted to that source. All specimens are housed in the T.M. Sperry Herbarium (KSP) at Pittsburg State University. A few have duplicates to be distributed.

RESULTS

In the following treatment, new records are listed alphabetically by taxon.

Ammannia robusta Heer & Regel (Lythraceae). Grand redstem. This native obligate wetland species is reported here for the first time from northeastern Oklahoma. It occurs elsewhere in 17 Oklahoma counties, and its broader range includes British Columbia and Ontario in Canada, interruptedly south to California, Texas, Louisiana, and east to Ohio, Kentucky and Tennessee (Kartesz 2024).

Voucher specimens: U.S.A. MISSOURI: CLAY CO.: Watkins Woolen Mill State Park, 39.400653, -94.261073, muddy clay area near lake edge, very abundant, 19 October 2024 (fr), *A. Fulkerson* 86 (KSP049008). **OKLAHOMA:** OSAGE CO.: Hulah Lake, 36.92822, -96.089839, sandy and rocky receding shoreline, 21 September 2024 (fr), *G. Gibbs* 34 (KSP047225). WASHINGTON CO.: Washington Cove of Copan Lake, 36.912295, -95.943798, receding shoreline of Copan Lake, sandy soil, 21 September 2024 (fl, fr), *G. McClain* 43 (KSP049368).

Andropogon virginicus L. Broom-sedge. This native species is widespread in the eastern United States.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075738, -94.613737, open grassy area amidst deciduous forest, occasional, 5 November 2023, *N. Snow 12108 & R. Mason* (KSP046689).

Arenaria serpyllifolia L. (Caryophyllaceae). Large thyme-leaf sandwort. This widespread nonnative species occurs across most of the U.S. and occupies most counties in Missouri and adjacent states.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0727004, -94.4297651, minimally disturbed meadow by gravel path, common, 30 May 2024, *R. Mason 146 & N. Snow* (KSP050146).

Bouteloua dactyloides (Nutt.) J.T. Columbus (Poaceae). Buffalograss. This is the first report of this widespread native species for Bates County, and it is somewhat east of its main geographical range in the Great Plains. The population was found in a low-lying area with numerous, robustly growing plants with female inflorescences. The landowner (W. Bradley, pers. comm. to N. Snow, 2024) believed that part of the property was never tilled.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, State Line Road, 38.085842, -94.612708, minimally disturbed, sometimes flooded, low-lying meadow with areas of open soil, common in one small area, 13 May 2024 (fr), *R. Mason 92 & N. Snow* (KSP050092).

Bouteloua hirsuta Lag. subsp. ***hirsuta*** (Poaceae). Hairy grama. The range of this native subspecies is mostly to the west and southwest. This is the first report for Lawrence County, Missouri, representing a range extension ca. 90 miles east from its nearest occurrence in Labette County, Kansas. The nearest Missouri collections are from Jackson County to the north, and Iron County to the east. It is listed as a Species of Conservation Concern (S2) in Missouri. This record may represent a naturalized occurrence from previous plantings, and may have originated from a seed contaminant in grass seed mixes.

Voucher specimen: U.S.A. MISSOURI: LAWRENCE CO.: Private property 7.26 km directly south of U.S. Hwy 174, 1.33 km East of K Hwy, 37.053114, -93.701591, in bottomland of disturbed prairie lowland, rare for this area, 5 October 2024 (fr), *G. Singer 6* (KSP047230).

Carduus nutans L. (Asteraceae). Musk thistle / Nodding plumeless thistle. Apart from Maine and Florida, this often aggressively invasive nonnative species is in all the contiguous states, where it often is declared noxious (e.g., in AR, KS, MO, and OK). It is known from most adjacent and nearby counties in Missouri and Kansas.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0734291, -94.4345963, common along disturbed grassy walking path, 30 May 2024, *R. Mason 126 & N. Snow* (KSP050126).

Carex granularis Muhl. ex Willd. (Cyperaceae). Limestone-meadow sedge. This native species occurs in all adjacent counties in Kansas and Missouri although it is considered rare in Kansas (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075739, -94.613737, disturbed roadsides adjacent to entry gate, rare, 13 May 2024, R. Mason 59 (KSP050059).

Carex microdonta Torr. & Hook. (Cyperaceae). Little-tooth sedge. Ranging from Arizona to Florida in the contiguous U.S., this native species occurs in seven other counties in Missouri, much of southeast Kansas, but becomes most widespread in central Oklahoma to the Hill Country and southeast Texas (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075974, -94.613475, open grassy area amidst deciduous forest, 21 April 2024, R. Mason 40 & N. Snow (KSP050040).

Carex sartwellii Dewey (Cyperaceae). Sartwell's sedge. The range of this native species is mostly confined to the northern half of the contiguous U.S. states. In Missouri it occurs in Jackson County to the north and three other counties on or near the Iowa border.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.085344, -94.611299, disturbed meadow on ridgeline in area of gravelly overburden and road track, common, 13 May 2024, R. Mason 100 & N. Snow (KSP050100).

Cerastium brachypetalum Desportes ex Pers. (Caryophyllaceae). Gray mouse-ear chickweed. In the U.S. this nonnative species has a sporadic distribution mostly in the southeast and is known from adjacent counties in Kansas and Missouri (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.082267, -94.608043, 21 April 2024, R. Mason 32 & N. Snow (KSP050032).

Croton glandulosus L. var. *septentrionalis* Muell.-Arg. (Euphorbiaceae). Vente connigo / Tooth-leaved croton. This native variety occurs in all adjacent counties except Miami County, Kansas (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.072094, -94.4322934, edge of forest in grassy clearing, occasional, 11 July 2024, R. Mason 179 (KSP050179).

Cyperus subsquarrosus (Muhl.) Bauter. (Cyperaceae). Small-flower halfchaff sedge. This native species, until recently included under *Lipocarpha micrantha* (Vahl) G.C. Tucker (Yatskievych 1999), is irregularly and not densely scattered mostly in the eastern half of the U.S., with some occurrences as far west as California and Oregon. In Missouri it is known nearby in Vernon and Bates counties.

Voucher specimen: U.S.A. KANSAS: BOURBON CO.: Elm Creek Lake, ca. 15 km SW of Fort Scott, 37.754427, -94.85421, occasional in wet soil, 7 October 2023, N. Snow 12102 (KSP046681).

Daucus pusillus Michx. (Apiaceae). Small wild carrot / American wild carrot. This native species occurs elsewhere in Kansas, to the west in Labette, Montgomery, Wilson, Chautauqua, Cowley counties and Leavenworth County to the north. It is more common in many counties of southwestern Missouri and areas south, east and west (Kartesz 2024).

Voucher specimen: U.S.A. KANSAS: CRAWFORD CO.: Crawford County Lake State Park, 37.64263, -94.81294, 14 Nov 2024 (fl, fr), J. Haworth 34 (KSP049361).

Dinebra panicea (Retz.) P.M. Peterson & N. Snow subsp. ***mucronata*** (Michx.) P.M. Peterson & N. Snow (Poaceae). Needle viper-grass / Red sprangletop. This native subspecies occurs from western New Mexico to the Ohio River region (Ohio and Kentucky), but is most common in the lower Mississippi and Missouri River drainages from Kansas and Missouri south to Louisiana and Texas (Snow 2003, Kartesz 2024), ranging south into Argentina (Snow 2012). It occurs in several adjacent counties in Oklahoma (e.g., Osage, Nowata and Rogers) and Kansas (Chautauqua and Montgomery).

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove of Copan Lake, 36.912295, -95.943798, sandy soil of receding shoreline, 21 September 2024 (fl, fr), G. McClain 40 (KSP047221).

Dinebra panicoides (J. Presl) P.M. Peterson & N. Snow (Poaceae). Amazon viper grass / Amazon sprangletop (Snow 2003). The range of this native species in the U.S. includes Texas and Oklahoma to Illinois, Virginia, and Georgia (Kartesz 2024). This report is the farthest west in Oklahoma, where it otherwise is known only from Ottawa, LeFlore and McCurtain counties.

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, common on sandy lakeshore, 17 September 2023 (fl), N. Snow 12096 et al. (KSP046676).

Diodia virginiana L. (Rubiaceae). Virginia buttonweed. The first report in Washington County for this native species; the nearest collections in Oklahoma are west in Osage and Pawnee counties, and east in Craig and Ottawa counties (Kartesz 2024).

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove of Copan Lake, 36.912295, -95.943798, sandy soil of receding shoreline, 21 September 2024 (fl, fr), G. McClain 46 (KSP049366).

Elaeagnus umbellata Thunb. (Elaeagnaceae). Autumn olive. The first report of this nonnative woody species for Bates County, which collectively with other counties, documents the species' northward and westward expansion (Kartesz 2024). Advocates of wildlife often promoted its planting for cover and forage.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.072094, -94.4322934, disturbed grassy meadow on forest edge, 8 August 2024 (fr), R. Mason 210 (KSP050210).

Echinochloa crus-galli (L.) P. Beauv. (Poaceae). Barnyard grass. The first report of this extremely widespread and weedy nonnative species that prefers seasonally and disturbed saturated soils. Although occurring in all fifty U.S. states and most of Canada, it surprisingly has not been vouchered yet (Kartesz 2024) for several nearby counties in Kansas (Bourbon, Linn) and Missouri (Henry, St. Clair). It is easily confused with the native species *Echinochloa muricata* (Stubbendieck et al. 2017) due to subtle differences in the spikelet morphology, so county distributions might be considered tentative until vouchers are confirmed.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0731024, -94.4313583, uncommon on gravel pond shore, 8 August 2024, R. Mason 233 (KSP050233).

Elephantopus carolinianus Raeusch. (Asteraceae). Carolina elephant's-foot. This record helps fill in the documented occurrences for this native species along the western-tier counties of Missouri. It is relatively widespread from eastern Texas to Illinois, Pennsylvania, and Florida.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0720515, -94.4315653, disturbed grassy meadow on forest edge, 24 August 2024 (fl), R. Mason 286 (KSP050286).

Eleusine indica (L.) Gaertn. (Poaceae). Goose grass / Indian goose-grass. This is the report of this widespread, nonnative species for Bates County. As a C4 grass, the North American distribution of this nonnative grass is more pronounced in the southern half of the U.S. It occurs in all surrounding counties except St. Clair County.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.072094, -94.4322934, forest edge in grassy clearing 11 July 2024 (fr) R. Mason 182 (KSP050182).

Elymus villosus Muhl. ex Willd. (Poaceae). Hairy wild-rye / Downy wild-rye. This native species occurs in many counties in eastern Kansas, most of Missouri, and areas mostly northward and eastward.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075738, -94.613737, open grassy area amidst deciduous forest, common, 5 November 2023, N. Snow 12106 & R. Mason (KSP046687).

Euonymus alatus (Thunb.) Sieb. (Celastraceae). Burning bush / Winged spindle-tree. This nonnative species is known nearby in Jasper (to the west) and Barry (to the south) counties, and several more counties mostly west, south and east within an approximately 100 km radius.

Voucher specimen: U.S.A. MISSOURI: LAWRENCE CO.: Private property 0.99 km east of K Hwy, 7.21 km south of U.S. Hwy 174 in Orange, 37.047304,

-93.703245, disturbed oak savanna on hill, 19 October 2024 (fr), G. Singer 20 (KSP047227).

Euphorbia prostrata Aiton. (Euphorbiaceae). Ground-fig spurge / Prostrate sand-mat. This record represents an eastern extension from Kansas where this native species is more common (Kartesz 2024), although it also occurs interruptedly to Massachusetts and Florida.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.073325, -94.431254, occasional on gavel shore of pond next to parking lot, 24 August 2024, R. Mason 274 (KSP050274).

Euploca convolvulacea Nutt. var. ***convolvulacea*** (Heliotropaceae). Seaside heliotrope. The first report for this native variety for Osage County; known from about 26 other counties in Oklahoma, mostly from the southern tier and western half of the state, with the nearest occurrence in Payne County, Oklahoma. The broader distribution of this native variety is mostly the southern tier states of the U.S. into the southern Great Plains, occurring as far north as Lancaster County, Nebraska.

Voucher specimen: U.S.A. OKLAHOMA: OSAGE CO.: Hulah Lake, vicinity of Skull Creek North Campground, 36.965046, -96.098154, sandy lakeshore, 29 September 2024 (fl), C. Helman 26 (KSP049360).

Fimbristylis autumnalis (L.) Roemer & J.A. Schultes (Cyperaceae). Slender fimbry. This native species occurs across most of the eastern U.S. and Canada with three outlying county occurrences in California (Kartesz 2024). In Oklahoma it is represented in over half of the eastern counties.

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, sandy lakeshore, 27 September 2024, K. Stringer 36 (KSP046712).

Fimbristylis vahlii (Lam.) Link (Cyperaceae). Vahl's fimbry. Ranging from California sporadically to Nebraska, Illinois and Virginia and Florida, the native species' range is most densely occupied in the southern Great Plains, east Texas, and the lower Mississippi River basin (Kartesz 2024).

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229; occasional on sandy lakeshore, 27 September 2024, K. Stringer 35 (KSP046711).

Gossypium hirsutum L. (Malvaceae). Upland cotton. The nearest other vouchered specimen is from Jasper County, Missouri. The species is grown widely in the southern third of the U.S. where it only infrequently becomes adventive beyond its native range in the southern third of Florida. The specimen has a large boll (fruit) and was the only specimen seen.

Voucher specimen: U.S.A. MISSOURI: LAWRENCE CO.: Private Property along north side of Farm Road 2000, 1.15 km East of U.S. Hwy 39, 2.65 km south of U.S. Hwy 60, 36.925299, -93.706631, along county road within disturbed prairie, 19 October 2024 (fr), G. Singer 17 (KSP047232).

Helianthus nuttallii Torr. & Gray subsp. ***rydbergii*** (Britt.) R.W. Long *vel aff.* Nuttall's sunflower. The identification here is tentative but based on several sources (Barkley 1986, Schilling 2006, SEINet 2025, Kartesz 2024). Hybridization and introgression are well known in sunflowers (Heiser 1969) and inevitably lead to some specimens that do not key easily or consistently using different keys, as was the case here. The specimen was collected from a population estimated at hundreds of stems from a rocky ledge (inaccurately indicated as "hill" on label). It keys imperfectly to the taxon indicated in *Flora of the Great Plains* (Barkley 1986) and *Flora of North America* (Schilling 2006). It differs from the description of the species in Schilling (2006) in having acute instead of 3-lobed chaff ("paleae") and lacking a pappus of 2 aristate scales. In other regards, however, the specimen compares well with four specimens of *H. nuttallii* in the Sperry Herbarium. Other comparisons were made with *H. x intermedius* R.W. Long, which is of hybrid origin (*H. maximiliani* Schrad x *grosseserratus* Martens) and according to BONAP (Kartesz 2024) is known from fewer than twenty counties from Kansas and Missouri to Maine. However, the specimen did not compare well to the one digitized image online (B.F. Bush 7779a, MO accession 809106). In addition, given over 30 digitized images (SEINet 2025) of the hybrid taxon *H. x kellermanii* Britt. (*H. grosseserratus* x *salicifolius* A. Dietr.), this possibility also was considered, but the leaf material is much narrower and longer on all those specimens compared to the one cited here. Leaf material is available upon request for DNA studies that might help better establish its identity or parentage.

Voucher specimens: U.S.A. KANSAS: CRAWFORD CO.: Bone Creek Lake, 37.621669, -94.745556, damp west-facing top of rocky hill next to muddy lake bank, 7 October 2023 (fl), I. McMahon 9 *et al.* (KSP046718 & KSP046719).

Heliotropium lasiocarpum Fisch. & C.A. Mey. L. (Heliotropiaceae). No evident common name yet for North America; suggestion based on its Latin binomial: Woolly-seed heliotrope. The native range of this nonnative, southern Eurasian species is from approximately Latitude 15–47°N and Longitude 25–115°E. Ours apparently is the second report of the species for North America and the first for the Great Plains and Oklahoma, the first being in 2005 from Maricopa County, Arizona (*J. Holt* 54 [ASU, image number ASU0092250]). Our specimen keys easily in *Flora Europea* to the *H. europaeum* "group," the four species of which the author (Brummitt 1972) indicated were not well understood and may not all merit recognition. Given that our specimen has a conical, papillate stigma and a moderate to dense indumentum on the smooth (not rugose) fruits, it keys easily to *H. lasiocarpum* Fisch. & C.A. Mey. Much more common in North America, as presently understood, is *H. europaeum* L., which occurs in over 100 counties scattered across the eastern U.S. (especially the Coastal Plain and Piedmont).

Voucher specimen: U.S.A. OKLAHOMA: OSAGE CO.: Hulah Lake, vicinity of Skull Creek North Campground, 36.965046, -96.098154, sandy shoreline, 29 September 2024 (fl, fr), G. Gibbs 18 (KSP047226).

Hemerocallis fulva (L.) L. (Asphodelaceae). Orange day-lily. The first report for Bates County. This nonnative, widely planted, vegetatively spreading species is known also from most adjacent counties to the north, west and south, but less documented in some counties to the southeast in Missouri and is widespread particularly in the eastern U.S.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.4235, -94.26072, 13 June 2024, uncommon in grassy roadside ditch, *R. Mason* 173 (KSP050173).

Hesperis matronalis L. (Brassicaceae). Dame's rocket. This nonnative species, planted frequently as an ornamental, has escaped widely across the U.S. and has been documented as adventive in most of the contiguous states (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.070847, -94.434825, disturbed grassland, common in scattered well-defined populations, 21 April 2024, *R. Mason & N. Snow* 48 (KSP050048).

Ipomoea purpurea (L.) Roth (Convolvulaceae). Common morning-glory. This nonnative species is designated as a noxious weed in Oklahoma and some surrounding states (Kansas, Arkansas). Collections are known from Kay and Washington counties to the west and east (respectively), with many additional counties in Oklahoma mostly in the eastern third of the state (Kartesz 2024). Reported also is its first occurrence in Bates County, Missouri, where it also is known from most adjacent counties to the west in Kansas and Cass County, Missouri, to the north, but not vouchered for many nearby counties east in Missouri.

Voucher specimens: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.072094, -94.4322934, disturbed grassy meadow on forest edge, 8 August 2024 (fl), *R. Mason* 203 (KSP050203). **OKLAHOMA:** OSAGE CO.: Hulah Lake, 36.92822, -96.089839, mixed sandy and rocky receding shoreline, 21 September 2024 (fl, fr), *G. McClain* 50 (KSP049359).

Lactuca floridana (L.) Gaertn. (Asteraceae). Woodland lettuce. The first record of this native species for Bates County; known from most surrounding counties in Missouri and Kansas and widespread in the eastern U.S. and parts of Manitoba and Ontario, Canada.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0734451, -94.4321475, uncommon in disturbed ditch on edge of concrete parking lot, 11 July 2024, *R. Mason* 183 (KSP050183).

Lonicera maackii (Rupr.) Maxim. Amur honeysuckle / Bush honeysuckle. This is an aggressively invasive and nonnative understory and ecotone shrub. Although it is known from the adjacent Vernon County to the south and approximately twelve other counties in the western half of Missouri, the species likely is more widespread in the general region than most county maps (e.g., Kartesz 2024) presently indicate. The private landowner was informed immediately of the small population of approximately ten plants.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075738, -94.613737, at base of escarpment, slightly upslope of seasonally saturated area of *Typha*, 5 November 2023, *N. Snow* 12124 & *R. Mason* (KSP046705).

Lysimachia nummularia L. (Primulaceae). Moneywort / Creeping jenny. This nonnative species is widespread in the U.S. but with most occurrences east of approximately Longitude 100°W. In Kansas it occurs in Cherokee, Neosho, and Bourbon counties to the south, west, and north (respectively) and approximately seventeen other counties in northcentral and eastern Kansas (Kartesz 2024).

Voucher specimen: U.S.A. KANSAS: CRAWFORD CO.: Mined Land Wildlife Area Unit #1, northeast corner of U.S. Hwy 160 and U.S. Hwy 69, 37.470045, -94.703048, in small seasonal pond, *A. Fulkerson* 50 (KSP049009).

Myosotis macrosperma Engelm. (Boraginaceae). Large-seed forget-me-not / Big-seeded scorpiongrass. Confirmation of this native species as a first report for Cherokee County was made by Douglas Goldman (USDA, pers. comm. to NS, Jan. 2025). In Kansas it is known only from Crawford County to the north and in Missouri it occurs in two adjacent counties (Jasper and Barton). Its broader range is mostly to the east and south. This specimen was collected for a MS thesis that later was published (Pryer et al. 2019).

Voucher specimen: U.S.A. KANSAS: CHEROKEE COUNTY.: On east side of NE 95th S ca. 85 m N of NE Bethlehem Rd., 37.19368, -94.65498, mixed deciduous forest and native tall grass prairie, 26 April 2014, *S. Pryer* 143 (KSP049513).

Oenothera biennis L. (Onagraceae). Common evening primrose / King's cure-all. This native species is known from all adjacent counties in Missouri and widespread across the eastern half of the U.S. and Canada, with fewer occurrences west of the Great Plains.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0727004, -94.4297651, uncommon on disturbed grassy slope next to rocky lakeshore, 8 August 2024, *R. Mason* 239 (KSP050239).

Oxalis corniculata L. (Oxalidaceae). Creeping yellow wood-sorrel. Widespread across all states in the contiguous U.S., this nonnative species is scattered across approximately 25 counties in Missouri, most of which are in the eastern third of the state.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, State Line Road, 38.082267, -94.608043, highly disturbed patchwork of secondary deciduous forest and grasslands, occasional, 21 April 2024, *R. Mason* 33 & *N. Snow* (KSP050033).

Papaver dubium L. (Papaveraceae). Blinneyes poppy. This nonnative ornamental species is known now in four counties in Kansas with widely scattered occurrences known from Missouri, Arkansas, Oklahoma, and about twenty other states. It was submitted to KSP by a local resident.

Voucher specimens: U.S.A. KANSAS: BOURBON CO.: Just inside Bourbon County line, near intersection of 100th Street and Old Picher Pike / Railroad Street, 37.680257, -94.942675, on roadside, 6 June 2023 (fr), *G. Eitel* s.n. (KSP046666).

OKLAHOMA: OTTAWA CO.: Lost Creek, along U.S. Highway 10, ca. 4.8 km NE of Wyandotte, 36.80294, -94.68878, locally common in grazed meadow/terrace above creek, 22 April 2023 (fl), *N. Snow* 12082 (KSP046663).

Paspalum laeve Michx. (Poaceae). Field crown grass / Field paspalum / Round seed paspalum. This native species is common mostly to the east and south and known from most counties in the southern two-thirds of Missouri.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075738, -94.613737, rare in open grassy area amidst deciduous forest, 5 November 2023, N. Snow 12120 & R. Mason (KSP046801).

Persicaria longiseta (Bruijn) Kitag. (Polygonaceae). Creeping smartweed. This nonnative species is widespread in the eastern half of the U.S., although relatively rare in western Missouri.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.073221, -94.431317, common on disturbed grassy slope next to rocky shoreline, 18 October 2024, R. Mason 414 (KSP050414).

Phlox paniculata L. (Polemoniaceae). Perennial phlox / Fall phlox. This widespread native species occurs in several Kansas counties, and is more common south and east in and beyond Missouri.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0705577, -94.4351958, disturbed ditch along gravel road, 8 August 2024 (fl), R. Mason 253 (KSP050253).

Phragmites australis (Cav.) Trin. ex Steud. (Poaceae). Common reed. Of the three congeners now recognized for North America, this one is nonnative and aggressively invasive in wetlands. It occurs across much of Canada and the 48 contiguous United States and is declared noxious in 15 of them. The specimen is sterile and less than 1 m tall but its strongly rhizomatous growth even in that early stage and other aspects of its morphology, coupled with its growth in a wetland situation, agree with species descriptions (e.g., lemma length and texture). The native *P. americanus* (Saltonst., P.M. Peterson & Soreng) A. Haines has a much less common distribution and SEINet indicates no occurrences for Oklahoma.

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, sandy lakeshore, only one plant seen, 27 September 2024, N. Snow 12090 (KSP046670).

Pinus strobus L. (Pinaceae). Eastern white pine. The collector observed this native species in a wooded area where an intentional planting seems unlikely, which if true represents another westward locality beyond its presumed pre-European contact North American distribution (Kartesz 2024). This collection is the farthest westward occurrence in Missouri, with the nearest collections being in Benton County, Arkansas and Crawford County, Missouri. Its native range is mostly north and east of Missouri, and this record may represent a naturalized occurrence from previous plantings.

Voucher specimen: U.S.A. MISSOURI: LAWRENCE CO.: Private property 7.38 km south of U.S. Hwy 174, 1.19 km East of K Hwy in Orange, 37.047683, -93.704209, in woodland, 21 September 2024 (female cone), G. Singer 13 (KSP047228).

Poa compressa L. (Poaceae). Flat-stem blue grass / Canada bluegrass. This nonnative species is widespread across the contiguous U.S. but is somewhat less common in the southeast (and as yet unknown for Florida [Kartesz 2024]).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0727004, -94.4297651, common in grassy clearing at forest edge, 30 May 2024, R. Mason 154 & N. Snow (KSP050154).

Podophyllum peltatum L. (Berberidaceae). Mayapple. This native species was known in all surrounding counties in Kansas and Missouri when discovered for the first time in Bates County in 2024. It is widespread across the eastern U.S.

Voucher specimen: MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.072167, -94.430407, many stems in patch at edge of deciduous forest, 19 May 2024, R. Mason 41 & N. Snow (KSP050041).

Polypremum procumbens L. (Tetrachondraceae). Juniperleaf. In the U.S. this native species is widespread from the mid-Atlantic states through the Gulf Coast to the southern tip of Texas and most of the southeast. In Oklahoma it occurs in about two dozen counties mostly in the eastern half of the state, the nearest being Delaware County.

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, sandy lakeshore, 27 September 2024. K. Stringer 37 (KSP046713).

Populus deltoides Bartr. ex Marsh. subsp. ***monilifera*** (Salicaceae). Eastern cottonwood. One of the most widespread native tree species in the continental U.S. This subspecies is largely restricted to the upper and central Great Plains, and while it is common throughout Kansas, its representation in Missouri is sparser.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.083484, -94.612146, occasional in disturbed meadows and adjacent gravelly roadsides near outbuildings, 13 May 2024, R. Mason 84 & N. Snow (KSP050084).

Potentilla rivalis Nutt. (Rosaceae). Brook cinquefoil. This native species previously was known from about fifteen counties in Missouri (Kartesz 2024) and is more common in states to the west.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.082267, -94.608043, highly disturbed patchwork of secondary deciduous forest and grasslands; only one plant seen, 21 April 2024 (fl), R. Mason 27 & N. Snow (KSP050027).

Prunus virginiana L. var. ***virginiana*** (Rosaceae). Choke-cherry. The typical variety of this widespread native species occurs mostly in the northern and central Great Plains and northern half of the eastern U.S., although it overlaps somewhat in the western part of its range with *P. virginiana* var. *demissa* (Nutt.) Torr. (Kartesz 2024). The typical variety occurs in other western tier counties in Missouri to the south and north of Bates County (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.082267, -94.608043, common in highly disturbed patchwork of secondary deciduous forest and grasslands, 21 April 2024 (fl), *R. Mason 21 & N. Snow* (KSP050021).

Quercus phellos L. (Fagaceae). Willow oak. Whereas this native species occurs in many counties in southeastern Missouri and across most of Arkansas (Kartesz 2024), this collection represents a northeastern extension from Delaware County, Oklahoma, and a northern extension from Washington County, Arkansas. Its occurrence at this site occasionally in the woodlands suggests a natural occurrence (G. Singer, pers. obsv.), and if so, then it is the westernmost station in Missouri. The specimens may represent a naturalized occurrence from previous plantings and, as noted by a reviewer, its relatively small acorns are readily moved by birds and mammals.

Voucher specimen: U.S.A. MISSOURI: LAWRENCE CO.: Private property 0.80 km north of U.S. Hwy 60, 0.23 km west of South Carnation Dr., 36.955334, -93.739387, drainage ditch leading into artificial pond and adjacent to disturbed prairie, 1 October 2024 (fr), *G. Singer 11* (KSP047229).

Ranunculus sardous Crantz (Ranunculaceae). Hairy buttercup. This nonnative species is more common south and east in North America (Kartesz 2024), but this location is the farthest north among the western-tier counties of Missouri.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Bradley Property, 9941 County Road 14367, 38.075739, -94.613737, infrequent on disturbed roadsides, 13 May 2024, *R. Mason 50 & N. Snow* (KSP050050).

Rorippa palustris (L.) Bess. subsp. ***palustris*** (Brassicaceae). Marsh yellowcress / Bog yellowcress. This native species is widespread across all North America including Greenland (Kartesz 2024).

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, sandy lakeshore, 27 September 2024 (fr), *N. Snow 12091* (KSP046671).

Sagittaria latifolia Willd. (Alismataceae). Duck-potato. This obligate native wetland species is common in Missouri, Kansas, and much of North America (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.073325, -94.431254, uncommon along gravel pond shore, 24 August 2024, *R. Mason 275* (KSP050275).

Sclerochloa dura (L.) Beauv. (Poaceae). Hardgrass / Common hardgrass. This nonnative cespitose (and typically prostrate) species now is widespread across much of North America, including British Columbia and Ontario, with at least thirty county occurrences combined in Kansas, Missouri, and Oklahoma (Kartesz 2024).

Voucher specimen: U.S.A. OKLAHOMA: OTTAWA CO.: Twin Bridges State Park, 36.798465, -94.753591, disturbed area of grassy parking lot, 22 April 2023. *N. Snow 12085* (KSP046660).

Sesbania herbacea (P. Mill.) McVaugh (Fabaceae). Peatree / Coffee weed. First report of this native species for Osage County. It has been documented in several counties nearby in Oklahoma (e.g., Washington, Nowata, Tulsa, Rogers) and Kansas (Lambette, Montgomery, Wilson) and becomes much more common mostly south and east (Kartesz 2024). It is declared a noxious weed in Arkansas and Louisiana.

Voucher specimens: U.S.A. OKLAHOMA: OSAGE CO.: Hulah Lake, 36.92822, -96.089839, common along receding rocky and sandy shoreline, 21 September 2024 (fl, fr [imm.]), S. Hulvey 28 (KSP049364). WASHINGTON CO.: Washington Cove on Copan Lake, 36.911033, -95.94229, sandy lakeshore, 27 September 2024. K. Stringer 44 (KSP046714).

Stellaria pallida (Dumort.) Crép. (Caryophyllaceae). Sand chickweed. Some authors include this nonnative species within *S. media* (L.) Vill. Although *Stellaria pallida* is not vouchered for the county, *S. media* is (Kartesz 2024).

Voucher specimen: U.S.A. OKLAHOMA: WASHINGTON CO.: Washington Cove of Copan Lake, 36.912295, -95.943798, sandy soil of receding shoreline, 21 September 2024 (fr), G. McClain 45 (KSP047222).

Strophostyles leiosperma (Torr. & A. Gray) Piper (Fabaceae). Slick-seed fuzzy-bean / Slick-seed wild bean. This native species is common in much of Missouri, Kansas and the central U.S. north to Ontario.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0734451, -94.4321475, disturbed ditch on edge of concrete parking lot, 8 August 2024 (fr), R. Mason 217 (KSP050217).

Triplidium ravennae (L.) H. Scholz (Poaceae). Ravenna grass, native to western Asia and southern Europe (Webster 2003) and widely planted as an ornamental, is published here for the first time for 20 counties in Kansas (see **Figure 1** for eastern Kansas records; also new to Crawford and Geary counties). It was previously known from nine counties in eastern Kansas (Styers 2022). In North America its distribution ranges from Washington to Michigan and New York, south to California and Florida, although it is largely absent from the central Great Plains and central and northern Rocky Mountains (Kartesz 2024, GBIF 2025). Species distribution modeling using MaxEnt (Styers 2022) indicated that most of eastern Kansas represents suitable habitat for the species. *Triplidium ravennae* also occurs along roadsides in several western-tier counties of Missouri but frequently cannot be collected given its presence on the margins of interstate highways, where roadside parking is prohibited. Synonyms that have been used in North America include *Erianthus ravennae* (L.) Beauv., *Erianthus ravennae* var. *purpurascens* (Anderss.) Hack., *Ripidium ravennae* (L.) Trin., and *Saccharum ravennae* (L.) L. Approximately half of the vouchers indicated below are photo vouchers, all deposited in the Sperry Herbarium and each with separate labels and barcodes. Sedgwick and Wilson counties each had fifteen collections.

Voucher specimens: U.S.A. KANSAS: ALLEN CO.: Iola, along northbound side of Hwy 169, ca. 1450 m SW of NW Minnesota Rd Exit, 13 September 2022 (fl), Styers 311 (KSP046554). BOURBON CO.: 2 km S of Prescott on northbound side

of Hwy 69, 28 August 2022 (fl), *Styers* 294 (KSP046537), *Styers* 296 (KSP046539), *Styers* 297 (KSP046540). BUTLER CO.: El Dorado, SW of Hwy 254 and SW Boyer Rd. Intersection, on NE corner of protected wetland, 13 September 2022 (fl), *Styers* 312 (KSP046555), *Styers* 313 (KSP046556), *Styers* 314 (KSP046557), *Styers* 315 (KSP046558), *Styers* 316 (KSP046559). CHAUTAUQUA CO.: Summit, 5 m N of westbound side of Hwy 166, ca. 1 km W of Rd. 14 intersection, 9 September 2022 (bud), *Styers* 304 (KSP046547). CHEROKEE CO.: Skidmore, Adjacent to Hwy 7 on E side, 15 April 2022 (fl), *Styers* 245 (KSP046489). CLOUD CO.: South of Concordia, ca. 20 m S of Hwy 81 and Airport Park Junction, 5 August 2022, *Styers* 271 (KSP046514), *Styers* 272 (KSP046515), *Styers* 273 (KSP046516), *Styers* 274 (KSP046517), *Styers* 275 (KSP046518). CRAWFORD CO.: Baker, S of E 530th Ave, 0.25 mi E of S 180th St., 1 May 2022 (sterile), *Styers* 247 (KSP046491), *Styers* 243 (KSP046487), *Styers* 246 (KSP046490), *Styers* 320 (KSP046563), *Styers* 321 (KSP046564), *Styers* 322 (KSP046565), *Styers* 323 (KSP046566), *Styers* 324 (KSP046567), *Styers* 325 (KSP046568), *Styers* 326 (KSP046569). DICKINSON CO.: West of Abilene, ca. 12 m N of westbound Hwy 70, 450 m E of Flag Rd. overpass, 5 August 2022 (bud), *Styers* 283 (KSP046526), *Styers* 284 (KSP046527), *Styers* 285 (KSP046528), *Styers* 305 (KSP046548), *Styers* 307 (KSP046550), *Styers* 308 (KSP046551), *Styers* 309 (KSP046552). FRANKLIN CO.: Hayes, adjacent to northbound exit to Stafford Rd., off Hwy 59, ca. 65 m NE of exit, 15 August 2022 (bud), *Styers* 288 (KSP046531). GEARY CO.: Junction City, ca. 50 m W of S Spring Valley Rd. overpass on eastbound Hwy 70, 5 August 2022, (fl), *Styers* 286 (KSP046529). GREENWOOD CO.: Salt Springs, adjacent to eastbound side of Hwy 400, ca. 265 m E of Aa Rd., 27 September 2022 (fl), *Styers* 336 (KSP046579). JOHNSON CO.: Overland Park, within exit loop of 63rd St. exit onto Metcalf Ave., 3 September 2022 (fl), *Styers* 301 (KSP046544). LABETTE CO.: Parsons, ca. 190 m E of Hwy 59 on S side of 22000 Rd., 7 August 2022 (fl), *Styers* 279 (KSP046522), *Styers* 280 (KSP046523), *Styers* 281 (KSP046524), *Styers* 282 (KSP046525). LINN CO.: Potosi, 4 mi N of Prescott on northbound side of Hwy 69, 28 August 2022 (bud), *Styers* 295 (KSP046538), *Styers* 298 (KSP046541), *Styers* 303 (KSP046546). LYON CO.: Emporia, 5 m N of roadside on Hwy 35 southbound, ca. 760 m E of Rd. R1 overpass, 4 August 2022 (bud), *Styers* 269 (KSP046512). MARSHALL CO.: Marysville, 4 m N of westbound edge of Hwy 36, 450 m W of 11th Rd., 5 August 2022 (bud), *Styers* 270 (KSP046513). MORRIS CO.: Herrington, adjacent to northbound side of Hwy 77, ca. 200 m S of E Helen St., 10 September 2022 (fl), *Styers* 306 (KSP046549), *Styers* 310 (KSP046553). OSAGE CO.: Fairfax, adjacent to southbound side of Hwy 75, ca. 650 m S of W 197th St. and Hwy 75 intersection, 15 August 2022 (bud), *Styers* 319 (KSP046562), *Snow* 11561 (KSP046664). SALINE CO.: Salina, ca. 35 m S of E Iron Ave., W of Smoky Hill River Bridge, 5 August 2022 (fl), *Styers* 276 (KSP046519), *Styers* 277 (KSP046520), *Styers* 278 (KSP046521). SEDGWICK CO.: Wichita, 4 m W of N Ridge Rd., off southbound edge, 1.1 mi S of Hwy 96 junction, 23 July 2022 (fl), *Styers* 249 (KSP046493), *Styers* 250 (KSP046494),

Styers 251 (KSP046495), Styers 252 (KSP046496), Styers 253 (KSP046497), Styers 254 (KSP046498), Styers 327 (KSP046570), Styers 328 (KSP046571), Styers 329 (KSP046572), Styers 330 (KSP046573), Styers 331 (KSP046574), Styers 332 (KSP046575), Styers 333 (KSP046576), Styers 334 (KSP046577), Styers 335 (KSP046578). WILSON CO.: New Albany, adjacent to Hwy 400 on N side, 1 April 2022 (fl), Styers 244 (KSP046488), Styers 248 (KSP046492), Styers 255 (KSP046499), Styers 256 (KSP046500), Styers 257 (KSP046501), Styers 258 (KSP046502), Styers 259 (KSP046503), Styers 260 (KSP046504), Styers 261 (KSP046505), Styers 262 (KSP046506), Styers 263 (KSP046507), Styers 264 (KSP046508), Styers 265 (KSP046509), Styers 266 (KSP046510), Styers 267 (KSP046511).

Typha latifolia L. (Typhaceae). Broad-leaf cat-tail. This native wetland species is common throughout most of North America (Kartesz 2024).

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.073325, -94.431254, uncommon on gravel shore of pond, 24 August 2024, R. Mason 277 (KSP050277).

Veronica missurica Raf. (Asteraceae). Missouri ironweed. This native species is more common in the eastern third and northern half of Missouri and occurs mostly west of the Mississippi and north of the Ohio rivers.

Voucher specimen: U.S.A. MISSOURI: BATES CO.: Harmony Mission Lake Conservation Area, 38.0739878, -94.4343894, highly disturbed grassy trail on edge of meadow, 8 August 2024 (fl), R. Mason 225 (KSP050225).

Vicia sativa L. subsp. *nigra* (L.) Ehrh. (Fabaceae). Garden vetch. This nonnative species occurs most commonly in the eastern U.S., although it is known in all 48 contiguous states. It occurs in about half of the counties in Oklahoma.

Voucher specimen: U.S.A. OKLAHOMA: OTTAWA CO.: Lost Creek, along U.S. Hwy 10, ca. 4.8 km NE of Wyandotte, 36.80294, -94.68878, locally common in grazed meadow/terrace above creek, 22 April 2023, N. Snow 12083 (KSP046662).




Figure 1. Distribution of *Tripidium ravennae* (L.) L. in eastern Kansas (from Styers 2022).

DISCUSSION

This paper summarizes recent collections or re-identifications of specimens occurring mostly within a 300 km radius of Pittsburg, Kansas. It includes one state record (*Heliotropium lasiocarpum*) from Oklahoma and 64 county records from Kansas, Missouri and Oklahoma. Given that first state or county occurrences sometimes are not traced easily in data aggregators such as SEINet (2025), print-based publications still represent an important source for reporting new records. Moreover, vouchered herbarium specimens likely will continue to be the preferred approach despite the advent of phone-based apps such as iNaturalist that capture digital images, especially given that digital sources frequently are unable to convincingly differentiate among taxa in species-rich genera (White et al. 2023, Eckert et al. 2024).

Previous publications of state records exclusively (Legler 2010) or state and county records together (Martine & Ward 2013, Snow et al. 2017, Pryer et al. 2019, Freeman & Morse 2019, Daines et al. 2022) have led to a growing realization that our knowledge of vascular plant distributions at the county level remains substantially incomplete. Such gaps in geographical

documentations are known as the Wallacean Shortfall (Brown & Lomolino 1998, Lomolino 2004, Snow et al. in press).

Whereas the primary focal level of new records previously was state records or significant range extensions (e.g., Daines & Snow 2024), digital tools such as SEINet and BONAP increasingly enable workers to determine quickly and with some degree of confidence whether a collection represents a first herbarium voucher for a state or county.

The plethora of records reported here and elsewhere is unsurprising in one sense, given that relatively few people track the distributions of plant species at the county level. Moreover, few outside of active and well-curated herbaria have the means to identify specimens of many taxa with a high degree of confidence, since well-curated comparative material is often critical for confirmation of identifications.

In contrast, it is surprising that so many county and state records are still being documented in areas such as Kansas, Missouri, and Oklahoma after (albeit sporadic) collecting for at least 140 years. This magnitude of county records may be underappreciated by many plant taxonomists and ecologists. As an example of an incompletely surveyed county, more specimens of vascular plants (ca. 415) were collected in Bates County, MO between November 2023 and April 2025 than from the 1970s through the 2010s (R. Mason & N. Snow, unpublished).

Another goal of plant taxonomy not mentioned in the Introduction is training the next generation of professionals in the proper techniques of field collecting and databasing. The first in-depth scientific exposure to plant diversity for most is through courses offered in colleges, universities, or workshops. Students collecting plants early in their careers, contributing to county and sometimes state records, and participating in the publication process provides critical experience and helps to instill the importance of local and current knowledge of plant distributions. Collections made by students from their counties of origin and on private properties will continue to be an important and ongoing source of novel plant distributions.

ACKNOWLEDGEMENTS

For permission to collect plants we thank William Bradley and Blackberry Farms in Missouri and Gabe McClain for facilitating that work; Malissa Briggler (Botanist, Missouri Department of Conservation); Brandon Moehrle (Lake Manager, U.S. Army Corps of Engineers, Tulsa District); and an anonymous landowner in Oklahoma. Thanks also to Doug Ladd and two anonymous reviewers for their close readings of the manuscript and suggested additions.

LITERATURE CITED

Barkley, T.M. 1986. Asteraceae. pp 838-1021 in: Great Plains Flora Association. *Flora of the Great Plains*. University Press of Kansas, Lawrence., pp. 952-957.
Brown, J.H. & M.V. Lomolino. 1998. *Biogeography*, 2nd edition. Sinauer Press, Sunderland, MA.

Brummitt, R.K. 1972. *Heliotropium*. pp. 84-86 in: Tutin, T.G., V.H. Heywood, N.A. Burges, D.M. Moore, D.H. Valentine, S.M. Walters, & D.A. Webb. *Flora Europaea*, volume 3. Diapensiaceae to Myoporaceae. Cambridge University Press, UK.

Daines, M. & N. Snow. 2024. Noteworthy Collections: Idaho. *Madroño* 71: 10–11.

Daines, M., A. Anzjon, M. Gay, L. Headings, J. Huckabee, M. Marine, T. Poolman, S. Scoggins, R. Styers, I. Villafaña, & N. Snow. 2022. Numerous distributional novelties collected by students for Idaho, Kansas, Missouri, Oklahoma, and Texas. *Journal of the Botanical Research Institute of Texas* 16: 551–557.

Eckert, I., A. Bruneau, D.A. Metsger, S. Joly, T.A. Dickinson, & L.J. Pollock. 2024. Herbarium collections remain essential in the age of community science. *Nature Communications* 15: 7586. <https://doi.org/10.1038/s41467-024-51899-1>.

Freeman, C.C., & C.A. Morse. 2019. New and interesting plant records for the central United States. *Missouriensis* 37: 16–38.

GBIF (Global Biodiversity Information Facility). 2025. *Tripidium ravennae* (L.) H. Scholz. <https://www.gbif.org/species/6431296>. Accessed 5 February 2025.

Haddock, M.J., C.C. Freeman, & J.E. Bare. 2015. *Kansas Wildflowers and Weeds*. University Press of Kansas, Lawrence.

Hammesfahr, A., P. Whitman, C. Campbell, & N. Snow. 2020. Another confirmation of *Cyperus flavescens* (Cyperaceae) for Kansas, U.S.A. *Journal of the Botanical Research Institute of Texas* 14: 411–412.

Heiser, C.B. 1969. The North American sunflowers (*Helianthus*). *Journal of the Torrey Botanical Club* 23: 1–218.

Kartesz, J. 2024. The Biota of North America's Floristic Synthesis of North America. Beta version dated 22 May 2024 (unpublished). University of North Carolina Botanical Garden, Chapel Hill.

Legler, B.S. 2010. Additions to the vascular flora of New Mexico. *Journal of the Botanical Research Institute of Texas* 4: 777–784.

Lomolino, M.V. 2004. Conservation biogeography. pp. 293-296 in: Lomolin, M.V. & L.R. Heaney, eds. *Frontiers of Biogeography: New Directions in the Geography of Nature*. Sinauer, Sunderland, MA.

Martine, C.T., & M.E. Ward. 2013. Establishment of regional herbarium leads to more than 200 new flora atlas records for New York. *Journal of the Torrey Botanical Society* 140: 125–131.

Pryer, S.Y., N. Snow, & J. Kartesz. 2019. Floristic survey of vascular plants in Crawford and Cherokee counties in southeastern Kansas. *Journal of the Botanical Research Institute of Texas* 13: 545–591.

Schilling, E.S. 2006. *Helianthus* L. pp. 141-169 in: Flora of North America Editorial Committee, eds. *Flora of North America North of Mexico*, volume 23: Magnoliophyta: Commelinidae (in part): Cyperaceae. Oxford University Press, New York and Oxford.

SEINet 2025. Southwestern Environmental Network Portal. <https://swbiodiversity.org/seinet>. Accessed 1 April 2025.

Snow, N. 2003. *Leptochloa* P. Beauv. pp 51-60 in: Flora of North America Editorial Committee, eds. *Flora of North America*, volume 25, Mangoliophyta: Commelinidae (in part): Poaceae, part 2. Oxford University Press, New York and Oxford.

Snow, N. 2012. *Leptochloa* P. Beauv. pp. 131-138 in: Zuloaga, F., Z. Rúgolo de Agrasar &, Z and A.R. Anton, eds. Flora Argentina: Flora Vascular de la República Argentina. Volumen 3. Tomo I. Monocotyledoneae. Poaceae: Aristidoideae a Pharoideae. Córdoba: Graficamente Ediciones.

Snow, N. 2017. New records of vascular plants for Kansas. *Phytoneuron* 2017-71: 1-2.

Snow, N., S. Young, C. Curran, & J. Kartesz. 2017. Vascular Plant Survey of the Neosho Bottoms Management Area (Grand River Dam Authority) in Ottawa County, Oklahoma. *Phytoneuron* 2017-26: 1-27. http://www.phytoneuron.net/2017Phytoneuron/26PhytoN_NeoshoFlora.pdf.

Snow, N., S. Young, & J. Kartesz. (In press). Reducing the Wallacean Shortfall: six decades of change in the vascular flora of Crawford County (Kansas, U.S.) highlights the importance of ongoing, targeted surveying. *Systematic Botany*.

Styers, R. 2022. Documenting the rapidly expanding distribution of Ravenna grass (*Tripsidium ravennae*) in eastern Kansas. M.S. thesis, Pittsburg State University, Pittsburg, Kansas.

Stubbendieck, J., S.L. Hatch, & C.D. Dunn. 2017. *Grasses of the Great Plains*. Texas A&M University Press, College Station.

Webster, R.D. 2003. *Saccharum* L. pp. 609-616 in: Flora of North America Editorial Committee, eds. Flora of North America, volume 25, Mangoliophyta: Commelinidae (in part): Poaceae, part 2. Oxford University Press, New York and Oxford.

White, E., P.S. Soltis, D.E. Soltis, & R. Guralnick. 2023. Quantifying error in occurrence data: Comparing the data quality of iNaturalist and digitized herbarium specimen data in flowering plant families of the southeastern United States. *PLOS ONE*. <https://doi.org/10.1371/journal.pone.0295298>.

Wieczorek J., D. Bloom, R. Guralnick, S. Blum, M. Döring, et al. 2012. Darwin Core: An Evolving Community-Developed Biodiversity Data Standard. *PLOS ONE* 7(1): e29715. <https://doi.org/10.1371/journal.pone.0029715>.

Yatskievych, G. 1999. Steyermark's Flora of Missouri, revised edition, volume 1. Missouri Department of Conservation, Jefferson City.

Yatskievych, G. 2006. Steyermark's Flora of Missouri, revised edition, volume 2. Missouri Botanical Garden Press, St. Louis.

Yatskievych, G. 2013. Steyermark's Flora of Missouri, revised edition, volume 3. Missouri Botanical Garden Press, St. Louis.